

# Maternity Nurses' Performance Regarding Cardiopulmonary Resuscitation During Pregnancy: Simulation Based Intervention

Aziza Ibrahim Mohamed<sup>1</sup>, Hemmat Mostafe Elbana<sup>1</sup>, Samah Abd Elhaleim<sup>1</sup>

<sup>1</sup> Maternal and Newborn Health Nursing, Benha University, Egypt

Correspondence: Aziza Ibrahim Mohamed Hassan, Assistant professor of Maternal and Newborn Health Nursing, Benha University, Egypt.

Received: March 5, 2018

Accepted: March 28, 2018

Online Published: April 2, 2018

doi:10.20849/ijsn.v3i1.356

URL: <https://doi.org/10.20849/ijsn.v3i1.356>

## Abstract

**Background:** Training maternity nurse regarding cardiopulmonary resuscitation during pregnancy is most important for properly care of such emergency situations during pregnancy and save woman's and fetal lives. The aim of the present study was to evaluate effect of simulation-based intervention on maternity nurse's performance regarding cardiopulmonary resuscitation during pregnancy. Design Aquasi-experimental design was adopted in the current study. The study was conducted at woman's health nursing department training lab at faculty of nursing affiliated at Benha University Hospital. Where a lot of women are admitted for normal and vaginal delivery, gynecological treatment and surgery.

**Subjects:** A Convenient sample of a total 52 maternity nurses was included.

**Tools:** Two tools: Were used for data collection. Interviewing questionnaire sheet was concerning with nurses personnel characteristics' and knowledge regarding cardiopulmonary resuscitation during pregnancy and performance check list sheet. The result of the present study that there was a highly statistical difference between knowledge and practical scores of maternity nurse at pre and post intervention. The study concluded that the simulation bases intervention is highly improved maternity nurses' performance regarding cardiopulmonary resuscitation during pregnancy. The present study recommended that simulation-based training regarding cardiopulmonary resuscitation should be provided for all obstetrics health care givers.

**Keywords:** simulation, maternity nurse, training program, performance, cardiac pulmonary resuscitation

## 1. Introduction

Cardiopulmonary arrest during pregnancy is life threatening condition and a leading cause of death. It presents a unique clinical scenario involving two patients: the mother and the fetus. The management of cardiac arrest in pregnancy is an important task for the emergency health team (Adam et al., 2016). The clinical outcome of mother or fetus will often depends on the successful resuscitation of the first few minutes. And a rapid multidisciplinary approach, including anesthesiology, medicine, obstetrics, neonatology, and sometimes cardiothoracic surgery. Basic and advanced cardiac life support algorithms should be implemented (Rittenberger, 2010).

The anatomic and physiologic changes during pregnancy result in significantly decreased cardiovascular and pulmonary reserves which complicate resuscitation. Cardiac output rises 30% to 50% as a result of increased stroke volume and to lesser extent increase in maternal heart Rate from 15-20bpm (Tan., 2013). Systemic vascular resistance decreased as a result of increase in several endogenous vasodilators including progesterone, estrogens and nitric oxide leading to decrease in mean arterial pressure, enlarge uterus can produce increased after load through compression of aorta and decrease cardiac return through the compression of inferior vena cava (Al-Foudri, 2010).

The estimates incidence of cardiac arrest during pregnancy ranged from 1/15,000 to 1/30,000-with maternal survival rates as low as 7% reported. The recently published scientific statement from the American Heart Association (AHA) addressing cardiac arrest in pregnancy is the first of its kind, even though a cardiac arrest risk in pregnancy has long been a known (Myerburg, 2015).

Cardiac arrest occurs only about one in every 30 000 late pregnancies, but survival from such an event is exceptional. Most deaths are from acute causes, with many mothers receiving some form of resuscitation.

However, the number of indirect deaths that is, deaths from medical conditions exacerbated by pregnancy is greater than that of deaths from conditions that arise from pregnancy itself (Micheal, 2014).

There are different causes of cardiac arrest during pregnancy; the most common cardiac cause is myocardial infarction, due to either premature atherosclerosis or coronary artery dissection. This relates in part to an increase in the proportion of pregnancies in women beyond the age of 35 years. The second most common cardiovascular cause is aortic aneurysms/dissections, and the third is the category of complex congenital heart diseases (Micheal, 2014).

The British center for maternal and child enquiries report has stated that timely recognition of pregnant women at risk of potentially life threatening conditions plays an important role in the appropriate institution of treatment (Wong et al, 2014).

Cardiopulmonary resuscitation during pregnancy is an integral part of health care of woman and fetus. Basic Life training (BLS) is highly essential for all healthcare staff members; especially to those who are working in obstetrics care units because protocol based management avoids confusion, wastage of time which save many lives. Survival after cardiopulmonary arrest is usually depends on early intervention, quality of cardiopulmonary resuscitation (CPR) and time to defibrillation (Demestihia et al., 2012).

Simulation is a technique for practice and learning that can be applied to many different disciplines and trainees; Simulation-based intervention is the way for developing health care provider knowledge, skills, and attitudes, in order to protect patients from unnecessary risks (Institute of Medicine, 2015).

The utilization of simulation to improve outcomes has been demonstrated in multiple domains because it gives the trainer the opportunity to experience uncommon scenarios, receive feedback, and perhaps most importantly correct mistakes before they become part of the learner's or groups mental model (Lavonas et al., 2015).

Simulation has also been shown to improve competencies in obstetric emergency decision-making, leadership, and individual and team performance This sentiment is reinforced in Committee Opinion Number 590 from The American College of Obstetricians and Gynecologists(ACOG), which states that one parameter for preparedness is integration of standardized emergency simulation and drills to identify and correct common clinical errors, to practice effective communication in a crisis, and to reinforce protocols, activation criteria, and critical interventions (Robertson et al., 2009).

Nurses are often first responders in cardiopulmonary arrest, and should be able to begin basic emergency care. Basic emergency care is crucial, rapid mobilization of expert resuscitation team and BLS performed competently until the arrival of these teams gives women the best chance for return of spontaneous circulation (Neal et al., 2012) Nurses play an important role in emergency management. Maternity nurse practice of resuscitation skills is best achieved through regular effective training. However, providing this teaching in a busy high-acuity maternity care units had become a challenge due to time and logistical constraints. A program to effectively and efficiently teach ICU nurses the skills necessary in patient resuscitation was developed using simulation training to better meet maternity learning needs (Jeejeebhoy et al., 2015).

### *1.1 Aim of the Study*

The aim of the present study was to evaluate effect simulation based intervention on maternity nurses' performance (knowledge and practice) regarding cardiopulmonary resuscitation during pregnancy.

### *1.2 Research Hypothesis*

Simulation based intervention will improve maternity nurses knowledge and practice regarding cardiopulmonary resuscitation during pregnancy.

### *1.3 Significance of the Study*

However cardiac arrest is rarely occurring during pregnancy. It is associated with a high maternal and fetal mortality rate. A multi-center study 2 reported the outcomes of 55 women who experienced cardiac arrest during pregnancy, including 12 who underwent perimortem cesarean delivery: only 15% (8/55) survived. So a survival rate of pregnant women is very poor.

In addition the center for maternal and child enquiries (CMACE) reports, concerning 2006-2008, have revealed that care was substandard in more than 50% of maternal deaths caused by cardiac arrest and that resuscitation skills were "considered very poor in an unacceptably high number of cases" (CMACE 2011). Several studies published on this matter (Cohen et al., 2008. Einav, et al., 2008) have shown a significant lack of knowledge among obstetric care providers about differences in the resuscitation of the pregnant women. A variety of medical professionals are

involved in the resuscitation of these women and, as cardiac arrest is such a rare occurrence on labor and delivery units, staff members are often unprepared to address this unexpected event.

So, providing high quality care quickly and efficiently to the pregnant patient who suffers cardiac arrest is a complex process, which requires previous awareness and training. From the clinical experience at the studied setting, the retention of cardiopulmonary resuscitation skills was poor. Particularly among maternity nurses who have little opportunity to practise CPR. Regular short periods of practice on a manikin are therefore essential. So the aim of the current was to evaluate effect of simulation based intervention on maternity nurses' performance.

## **2. Subject and Methods**

### *2.1 Research Design*

A quasi-experimental design was used in the present study.

### *2.2 Research Setting*

Woman's health nursing department training lab at faculty of nursing affiliated at Benha university hospital.

### *2.3 Subjects*

A convenient sample of a total 52 maternity nurses who working at obstetric and gynecological department affiliated at Benha university hospital.

### *2.4 Tools of Data Collection: Two Tools Was Used to Collect Data*

Tool I-Self-administered interview questionnaire this tool consisted of 2 parts:

Part 1: include the nurse's demographic data such as age, educational level, educational qualification, years of experience.

Part 2: to assess maternity nurse knowledge regarding cardiac arrest during pregnancy and cardiopulmonary resuscitation. It includes 16 different questions. Each questions given a correct answer, the total number of correct answers were 59 answers, answer was scored as (2) for correct answer and (1) for incorrect answer. The total knowledge score was calculated by summation of the scores for the correct answers. The total possible score ranged from (1 to 118 marks) and means and standard deviations were calculated. The higher scores reflect higher levels of knowledge about CPR during pregnancy.

The total knowledge score was indicated as the following:

Total knowledge score was calculated as:

Adequate knowledge  $\geq 60\%$  of total knowledge score ranged from (71-118 marks).

Inadequate knowledge  $< 60\%$  of total knowledge score ranged from (1-70 marks)

Tool II-maternity nurse practice observational checklist

It was constructed by the researcher after reviewing relevant literature. It was adopted from (Hui et al., 2011, Rodrigues et al., 2014), and (Adams et al., 2016). It was concerned with assessing the maternity nurse practice regarding cardiopulmonary resuscitation during pregnancy. This part consisted of (66 steps) items that identify the steps of different cardiopulmonary resuscitation. Preparatory and basic intervention (12 steps), compression related practice (8steps), open air way and breathing technique include (15 steps), defibrillation technique (17 steps), and intubation and post resuscitation care related practice included (14 steps). Each step was scored as (1) for not done, and (2) for done, total practice score as satisfactory practice  $\geq 75\%$  of total practice score (99-132marks), unsatisfactory practice  $< 75\%$  (1-99 marks).

### *2.5 Pilot Study*

A pilot study was conducted on (10%) of the studied nurses to test the feasibility, clarity and applicability of the developed tools, based on the finding of the pilot study.

### *2.6 Tool Validity*

Tools developed by the researchers were examined by a panel of five experts (three cardiologists and two professors of maternity nursing speciality) to determine whether the included items clearly and adequately cover the domain of content addressed. The percentage of consensus among experts regarding the structured interviewing questionnaire was 94% and the pre-post-test was 87%.

### 2.6.1 Tool Reliability

Test-retest was repeated to the same sample of maternity nurses on two occasions and then compares the scores. The Cronbach's coefficient alpha was 0.879.

### 2.7 Simulation Based Intervention

The simulation intervention provided by the researchers using different strategies, and training sessions in individual basis, small and large group discussion strategies also were utilized. Simulation training was conducted at the woman's health nursing department training lab affiliated at faculty of nursing at Benha University. Researchers prepare the training lab with essential needed supplies as Gaumard's NOELLE maternal birthing simulator with a pregnant uterus was used for all simulations to enable the performance of breathing, chest compression, intubation. And defibrillation in addition a digital monitor displaying dynamic vital signs was readily to evaluate woman's condition and assessment fetal wellbeing device.

#### 2.7.1 Ethical Consideration

An Informed consent was obtained from each maternity nurse at the first session. Each maternity nurse was informed that the participation in the study was voluntary and can withdraw at any time.

#### 2.7.2 Procedure

The study was begun at the 1st of February 2015 after receiving the approval from the dean of faculty of nursing and the director of Benha university hospital to the end of July 2015. Maternity nurses were divided into 10 different small groups in order to avoid contradiction with their work, each group contain 5 to 6 nurses. Simulation intervention was implemented through 3 different sessions (one theoretical session and two practical training sessions) for each group the researchers provide firstly a session regarding theoretical knowledge about cardiac arrest and cardiopulmonary resuscitation during pregnancy (meaning, causes, factors aggravate cardiac arrest during pregnancy) at the practical sessions were concerned with training nurses regarding technique of cardiopulmonary resuscitation during pregnancy including breathing technique, chest compression technique, defibrillation and intubation technique.

#### 2.7.3 Evaluations

The pre-posttest format was used in two times of assessment; the first was before the intervention, the second time was after the implementation of the simulation intervention which extends to three months. For evaluating practical skills maternity nurses were asked to demonstrate the procedure on the studied simulator for three different times and average of different score was calculated.

### 2.8 Statistical Design

Data analysis was performed using IBM SPSS (Statistical Package for Social Sciences) statistical software version 22. The data were explored. Descriptive statistics with mean and standard deviation (SD) for continuous variables and frequency for categorical variables were analyzed. Qualitative variables were compared using chi square test ( $\chi^2$ ) as the test of significance, paired and independent (t) test and ANOVA test was used to compare mean score between two and more groups respectively. Correlation coefficient (r) was used to evaluate association between studied variables. The p-value is the degree of significant. A significant level value was considered when  $p\text{-value} \leq 0.05$  and a highly significant level value was considered when  $p\text{-value} \leq 0.001$ , while  $p\text{-value} > 0.05$  indicates non-significant results.

## 3. Results

Table 1: Reveals that more than half of the studied maternity nurses were aged (20-29) years, more than half (54.0%) of them had a secondary nursing education. Moreover 38.0% of them had less than five years of experience. In addition more than half live in urban area. It also indicates that the highest percentage 78% previous training regarding CPR.

Table 2: Shows that, there was a highly statistically significant difference between mean scores of knowledge related to cardiac arrest during pregnancy between the pre and post intervention phases ( $p < 0.001$ ).

Figure 1: Percentage distribution of total knowledge score of the studied nurses at different phases of the program.

Table 3: Reveals that, there was a highly statistically significance difference between total knowledge scores of maternity nurses and their educational qualification, years of experience and previous training regarding CPR.

Table 4: Indicates that, there was a highly statistically significant difference between studied nurses' practice about basic intervention for cardiopulmonary resuscitation procedure between the pre and post intervention phases ( $p < 0.001$ ).

Table 5: Indicates that, there was a highly statistically significant difference between studied nurses' practice related to applying chest compressions between the pre and post intervention phases ( $p < 0.001$ ).

Table 6: Indicates that, there was a highly statistically significant difference between studied nurses' practice related to open airway and delivering breath technique at both the pre and post intervention phases ( $p < 0.001$ ).

Table 7: Shows that, there was a highly statistically significant difference between studied nurses' practice regarding defibrillator technique between the pre and post intervention phases ( $p < 0.001$ ).

Table 8: Indicates that, there was a highly statistically significant difference between studied nurses' practice scores related to applying intubation between the pre and post intervention phases ( $p < 0.001$ ).

Figure 2: Percentage distribution of total practice score of the studied nurses at different phases of the program.

Table 9: Reveals that, there was a highly statistically significance difference between total practice scores of maternity nurses and their previous training regarding CPR, at post intervention phase ( $p < 0.001$ ) and, moreover there was statistically significance difference between total practice scores in relation to their educational qualification at post intervention phase ( $< 0.05^*$ ).

Table 10: Shows that, that there was a highly positive association between studied nurses knowledge and practice scores at post intervention phase.

Table 1. Distribution of personnel characteristics of the studied nurses

Variable	Frequency	%
<b>Age in years</b>		
20-29	26	52.0
30-39	20	40.0
40-50	4	8.0
<b>Mean <math>\pm</math>SD</b>	28.23 $\pm$ 5.02	
<b>Educational qualification</b>		
Secondary nursing education	27	54.0
Technical nursing education	18	36.0
Baculare of nursing	5	10.0
<b>Years of experience</b>		
5-9	19	38.0
10-14	8	16.0
15-19	11	22.0
19-25	12	24.0
<b>Mean <math>\pm</math>SD</b>	10.58 $\pm$ 6.35	
<b>Residence</b>		
Urban	32	64.0
Rural	18	36.0
<b>Previous training regarding CPR</b>		
No	39	78
Yes	11	22

Table 2. Distribution of knowledge regarding cardiac arrest during pregnancy of the studied nurses at different phases of the intervention

Knowledge	No of answers	Pre-intervention Mean ± SD	Post-intervention Mean ± SD	Paired t test	P value
Definition of cardiac pulmonary arrest	2	1.3200±.47121	1.7600±.43142	-4.831	<0.001**
Causes of cardiac pulmonary arrest in pregnancy	6	1.7800±1.20017	4.9000±1.75255	-10.995	<0.001**
Complication of cardiac arrest during pregnancy	5	1.7400±.80331	3.7000±1.47427	-8.031	<0.001**
Management of cardio pulmonary during pregnancy	5	1.5600±.86094	4.1800±1.40973	-9.837	<0.001**
<b>Cardiopulmonary resuscitation :</b>					
Definition of CPR	2	1.2400±.43142	1.8400±.37033	-7.425	<0.001**
Importance of CPR	6	1.9800±.76904	4.9800±1.65973	-11.813	<0.001**
General principles of CPR	4	1.6400±.48487	3.3600±.96384	-11.369	<0.001**
Indication for CPR	3	1.3800±.49031	2.8400±.42185	-15.992	<0.001**
Ways to open the airway	3	1.2600±.44309	2.6200±.66670	-11.627	<0.001**
Depth of chest compression	3	1.3200±.5107	2.8200±.43753	-13.908	<0.001**
Rate of compressions	2	1.2800±.45356	1.6600±.47852	-4.030	<0.001**
Rate of chest compression to the breathing	2	1.3000±.46291	1.9400±.23990	-8.041	<0.001**
Site at which starts chest compressions	2	1.3400±.47852	1.7800±.41845	-4.831	<0.001**
Complications of CPR	5	1.7400±.82833	4.5600±1.10951	-15.667	<0.001**
Physiological changes during pregnancy affect cardio pulmonary resuscitation.	5	1.4000±.69985	4.5000±1.16496	-14.467	<0.001**
Drugs used during cardio pulmonary resuscitation during pregnancy	4	1.3200±.47121	3.7400±.63278	-21.116	<0.001**
Total	59	23.6000±4.00510	51.1800±3.91590	-35.411	<0.001**




Figure 1. Percentage distribution of total knowledge score of the studied nurses at different phases of the program

Table 3. Relation between studied nurses' total knowledge score and their personnel characteristics at different phases of intervention

Variable	Pre-intervention			Post-intervention				
	Mean ±SD	F test	Independent t test	P value	Mean ±SD	F test	Independent t test	P value
<b>Age in years</b>		1.34		>0.05		0.670		>0.05
20-	22.9231±4.61236				51.1923±4.08920			
30-	23.9500±3.06894				50.7500±3.87808			
≥ 40	26.2500±3.30404				53.2500±2.98608			
<b>Educational qualification</b>		10.49		<0.001**		8.74		<0.001**
Secondary nursing education	22.9630±2.91523				50.7037±3.17217			
Technical nursing education	22.7222±4.34952				50.2222±4.09527			
Baculare of nursing	30.2000±.44721				57.2000±.44721			
<b>Years of experience</b>		39.72		<0.001**		38.14		<0.001**
5-9	20.2609±2.54447				47.9130±2.77837			
10-14	23.0000±.81650				50.7500±.50000			
15-19	25.7273±1.48936				53.2727±1.27208			
≥20	28.2500±2.17945				55.6667±1.61433			
<b>Residence</b>		1.35		>0.05			0.227	>0.05
Rural	22.9688±3.30673				51.0938±4.36525			
Urban	24.7222±4.92061				51.3333±3.06786			
<b>Previous training regarding CPR</b>			3.09	<0.001**			2.88	<0.001**
No	22.7949±3.83323				50.4872±3.89920			
Yes	26.4545±3.35749				53.6364±2.97566			

Table 4. Distribution of practice regarding cardiopulmonary resuscitation procedure among the studied nurses at different phases of intervention

Basic intervention	Pre-intervention				Post-intervention				Chi-square $\chi^2$	P Value
	Not Done		Done		Not Done		Done			
	No	%	No	%	No	%	No	%		
<b>Pre procedure:</b>										
Activate maternal cardiac arrest team	45	90.0	5	10.0	6	12.0	44	88.0	60.86	<0.001**
Document time of onset of maternal cardiac arrest	44	88.0	6	12.0%	6	12.0	44	88.0	57.76	<0.001**
Keep women at left lateral position	43	86.0	7	14.0	9	18.0	41	82.0	46.31	<0.001**
Estimate gestational age	43	86.0	7	14.0	7	14.0	43	86.0	51.84	<0.001**
Prepare equipment	46	92.0	4	8.0	7	14.0	43	86.0	61.06	<0.001**
Keep privacy	39	78.0	11	22.0	7	14.0	43	86.0	41.22	<0.001**
<b>During procedure (Basic intervention):</b>										
Assess woman's consciousness	41	82.0	9	18.0	6	12.0	44	88.0	49.17	<0.001**
Don't transport women	41	82.0	9	18.0	9	18.0	41	82.0	40.96	<0.001**
Immediately call out for help	43	86.0	7	14.0	6	12.0	44	88.0	54.78	<0.001**
Palpate the carotid artery	43	86.0	7	14.0	7	14.0	43	86.0	51.81	<0.001**
Ensure a clear air way	45	90.0	5	10.0	6	12.0	44	88.0	60.86	<0.001**
Assess fetal heart sound	41	82.0	9	18.0	6	12.0	44	88.0	49.17	<0.001**

Table 5. Distribution of practice about demonstration of chest compression technique among the studied nurses at different phases of the program

applying Chest compressions	Pre-intervention				Post-intervention				Chi-square $\chi^2$	P Value
	Not Done		Done		Not Done		Done			
	No	%	No	%	No	%	No	%		
Hand position	42	84.0	8	16.0%	6	12.0	44	88.0	51.92	<0.001**
Place women at supine position during compression	44	88.0	6	12.0	3	6.0	47	94.0	67.48	<0.001**
Place hand in the centre of chest, slightly above the sternum	39	78.0	11	22.0	6	12.0	44	88.0	44.00	<0.001**
Chest compressions should be performed at a rate of at least 100 per minute.	46	92.0	4	8.0	9	18.0	41	82.0	55.31	<0.001**
Chest compressions should be performed at a depth of at least 2inch (5 cm)	40	80.0	10	20.0	6	12.0	44	88.0	46.53	<0.001**
Allow complete chest recoil after each chest compression	40	80.0	10	20.0	12	24.0	38	76.0	31.41	<0.001**
Minimize interruption	38	76.0	12	24.0	8	16.0	42	84.0	36.23	<0.001**
Perfume continuous manual left uterine displacement	41	82.0	9	18.0	11	22.0	39	78.0	36.05	<0.001**

Table 6. Distribution of practice of open air way and delivering breath technique among the studied nurses at different phases of the intervention

Delivering breath	Pre-intervention				Post-intervention				Chi-square $\chi^2$	P Value
	Not Done		Done		Not Done		Done			
	No	%	No	%	No	%	No	%		
<b>Open the airway and give breathing:</b>										
<b>Head-tilt/chin-lift manoeuvre</b>										
Hand position	44	88.0	6	12.0	10	20.0	40	80.0	46.53	<0.001**
Place the fingers of the other hand under the bony part	42	84.0	8	16.0	9	18.0	41	82.0	43.57	<0.001**
Grasp the angles of the woman 's lower jaw and lifting with both hands	46	92.0	4	8.0	7	14.0	43	86.0	61.06	<0.001**
Ensure a clear air way	43	86.0	7	14.0	6	12.0	44	88.0	54.78	<0.001**
<b>Deliver breathing</b>										
<b>A- perform the bag valve mask</b>										
Ensure a tight seal between the mask and the woman's face	46	92.0	4	8.0	6	12.0	44	88.0	64.10	<0.001**
Squeeze the bag with two hand	46	92.0	4	8.0	7	14.0	43	86.0	61.06	<0.001**
Administer 100% oxygen at 15 litre/minute	38	76.0	12	24.0	11	22.0	39	78.0	29.17	<0.001**
Compress the ambo-bag in rhythmical fashion	39	78.0	11	22.0	9	18.0	41	82.0	35.06	<0.001**
Attach the bag to an oxygen source	38	76.0	12	24.0	10	20.0	40	80.0	31.41	<0.001**
Pinch the woman's nostrils closed	40	80.0	10	20.0	7	14.0	43	86.0	43.71	<0.001**
Put the mouth completely over the woman's mouth	44	88.0	6	12.0	7	14.0	43	86.0	54.78	<0.001**
After 30 chest compression, give 2 breaths	45	90.0	5	10.0	9	18.0	41	82.0	52.17	<0.001**
Deliver each rescue breath over one second with enough force	47	94.0	3	6.0	10	20.0	40	80.0	55.85	<0.001**
Give sufficient tidal volume to produce visible chest rise	45	90.0	5	10.0	6	12.0	44	88.0	60.86	<0.001**
Avoid excessive ventilation	47	94.0	3	6.0	12	24.0	38	76.0	50.64	<0.001**



Table 7. Distribution of practice of defibrillator technique among the studied at different phases of the intervention

Basic intervention	Pre-intervention				Post-intervention				Chi-square $\chi^2$	P Value
	Not Done		Done		Not Done		Done			
	No	%	No	%	No	%	No	%		
Verify that woman is unresponsive.	42	84.0	8	16.0	10	20.0	40	80.0	41.02	<0.001**
Verify ECG reading of ventricular tachycardia, or ventricular fibrillation	40	80.0	10	20.0	10	20.0	40	80.0	36.00	<0.001**
Plug defibrillator into electric outlet.	39	78.0	11	22.0	13	26.0	37	74.0	27.08	<0.001**
Turn on defibrillator power.	41	82.0	9	18.0	9	18.0	41	82.0	40.96	<0.001**
Prepare woman and/or paddles with proper conductive agent	37	74.0	13	26.0	10	20.0	40	80.0	29.26	<0.001**
Checks that defibrillator is in asynchronous mode.	43	86.0	7	14.0	9	18.0	41	82.0	46.31	<0.001**
Deliver defibrillated										
Turn on ECG recorder.	42	84.0	8	16.0	9	18.0	41	82.0	43.57	<0.001**
Command all persons to move away from bed	40	80.0	10	20.0	8	16.0	42	84.0	41.02	<0.001**
Turn off oxygen source during actual	40	80.0	10	20.0	10	20.0	40	80.0	36.00	<0.001**
Stand away from bed area	41	82.0	9	18.0	11	22.0	39	78.0	36.05	<0.001**
Places the paddles at proper site on chest.	38	76.0	12	24.0	10	20.0	40	80.0	31.41	<0.001**
Apply paddles with firm pressure(11kg pressure)	37	74.0	13	26.0	10	20.0	40	80.0	29.26	<0.001**
Depress discharge buttons on defibrillator.	42	84.0	8	16.0	12	24.0	38	76.0	36.23	<0.001**
Deliver 3 stacked shocks, with biphasic shock energy of 120 to 200	38	76.0	12	24.0	13	26.0	37	74.0	25.01	<0.001**
Compressions should be resumed immediately delivery of the electric shock	36	72.0	14	28.0	13	26.0	37	74.0	21.16	<0.001**
Assess women cardiac rhythm	35	70.0	15	30.0	10	20.0	40	80.0	25.25	<0.001**
If first defibrillation unsuccessful, immediately charge paddles to 300 joules	33	66.0	17	34.0	15	30.0	35	70.0	12.98	<0.001**

Table 8. Distribution of practice of intubation technique among the studied nurses at different phases of intervention

Application of intubation	Pre-intervention				Post-intervention				Chi-square $\chi^2$	P Value
	Not Done		Done		Not Done		Done			
	No	%	No	%	No	%	No	%		
<b>Applying intubation:</b>										<0.001**
Continue to ventilation & oxygenation	40	80.0	10	20.0	7	14.0	43	86.0	43.71	<0.001**
Prepare Equipment	47	94.0	3	6.0	6	12.0	44	88.0	67.48	<0.001**
The anaesthetist may require	46	92.0	4	8.0	6	12.0	44	88.0	64.10	<0.001**
Recommence ventilation and oxygenation once intubation is completed	42	84.0	8	16.0	5	10.0	45	90.0	54.95	<0.001**
Continues CPR once intubation	43	86.0	7	14.0	8	16.0	42	84.0	49.02	<0.001**
Established venous access	44	88.0	6	12.0	8	16.0	42	84.0	51.92	<0.001**
Asepsis should be maintaining	43	86.0	7	14.0	7	14.0	43	86.0	51.84	<0.001**
The correct rate of infusion is required	46	92.0	4	8.0	7	14.0	43	86.0	61.06	<0.001**
Accurate documentation	45	90.0	5	10.0	7	14.0	43	86.0	57.85	<0.001**
<b>Applying Post resuscitation care:</b>										
Assess woman 's condition	45	90.0	5	10.0	8	16.0%	42	84.0	54.95	<0.001**
Check arterial blood gases	43	86.0	7	14.0	8	16.0%	42	84.0	49.02	<0.001**
Monitor woman's cardiac rhythm	39	78.0	11	22.0	9	18.0%	41	82.0	36.05	<0.001**
Continue respiratory therapy	36	72.0	14	28.0	7	14.0%	43	86.0	34.31	<0.001**
Assess woman's level of consciousness	38	76.0	12	24.0	5	10.0%	45	90.0	44.43	<0.001**




Figure 2. Percentage distribution of total practice score of the studied nurses regarding CPR during pregnancy at different phases of intervention

Table 9. Relation between studied nurses total practice score and their personnel characteristics at different phases of intervention

Variable	Pre-intervention				Post-intervention		
	Mean ±SD	F test	Independent t test	P value	Mean ±SD	F test	P value
<b>Age in years</b>		0.530		>0.05		1.72	>0.05
20-	73.8462±11.73607				118.8462±9.455		
30-	77.7500±15.36529				120.9500±8.331		
40-52	77.7500±12.97112				111.7500±10.59		
<b>Educational qualification</b>		57.69		>0.05		13.71	<0.05*
Secondary nursing education	69.1852±5.14435				114.2963±9.063		
Technical nursing education	76.7778±7.53467				123.3333±5.434		
Baculare of nursing	107.2000±14.5155				130.0000±0.000		
<b>Years of experience</b>		27.57		>0.05		0.258	>0.05
5-	70.2632±4.60485				116.4737±8.342		
10-	67.2500±3.24037				110.1250±7.510		
15-	71.0000±4.89898				118.5455±6.055		
20-24	94.3333±14.72989				129.8333±3.389		
<b>Residence</b>		0.202		>0.05		0.542	>0.05
Rural	76.0313±11.89872				119.6563±9.265		
Urban	75.1667±15.76761				118.1667±9.363		
<b>Previous training regarding CPR</b>		2.11		>0.05			<0.001**
No	72.7692±8.65558				117.7692±9.609	2.61	
Yes	86.1818±20.57580				123.9091±5.872		

Table 10. Correlation between studied nurses' knowledge and practice among studied nurses

Variable	Practice pre		Practice post	
	r	P value	r	P value
<b>Knowledge pre</b>	-.186	.195	-	-
<b>Knowledge post</b>	--	--	.485**	.000

\*\* . Correlation is significant at the 0.01 level (2-tailed).

#### 4. Discussion

Nurses are usually the primary witnesses to in-hospital sudden cardiac arrests. Training of maternity nurses has an impact on the efficiency and outcome of cardiopulmonary resuscitation (CPR) during pregnancy. So, the aim of present study was to evaluate the effect of simulation based educational intervention concerning CPR during pregnancy among maternity nurses. The present study findings approved that the utilization of a simulation based educational intervention concerning cardiopulmonary resuscitation (CPR) during pregnancy increase the knowledge and skills of maternity nurses and motivated them to comply with training.

Regarding the socio-demographic characteristics of the maternity nurses, the present findings showed that studied nurses age ranged from 20-50 years, and about one half of studied maternity nurses had a age group ranged from 20 to 25 years with years of experience ranged from 5-9 with mean of (10.58±6.35), and more than one half of them had a secondary nursing education. These study findings are agreed with Hussain et al. (2009) in the study at Bahrain, to evaluate nurses knowledge regarding cardiopulmonary resuscitation. It was reported that studied nurses' age ranged from 20-50 years and the majority had secondary nursing education, in addition, average years of experience was 12 years. Also (Mustafa & Jaddoue) in the study at Al-Najaf city's teaching hospital to evaluate Knowledge of nurses towards cardiopulmonary resuscitation.

In addition, the result of the current study showed that the majority of nurses hadn't previous training course related CPR during pregnancy. This result is in accordance with Damjan et al. (2012). in the study to evaluate effects of using of the impedance threshold device in cardiopulmonary resuscitation. Who stated that more than third of the nurses don't have any previous training regarding CPR, and mostly all of them were willing to attend training program. On the other hand, Nagashema et al. (2012) in the study to investigate nurses' knowledge and experience in cardiopulmonary resuscitation, was in contrast with the present study and stated that the majority of the nurses are much interested in CPR, and most of them had received education and training in CPR as students or after the graduation.

Additionally, the finding of the present study proved majority of studied maternity nurses' had inadequate level of knowledge regarding all knowledge items concerning CPR during pregnancy at pre intervention phase. These findings may be due to that there is lack of implementing educational training program regarding cardiopulmonary resuscitation for pregnant women at the studied setting. These findings are agreed with (Cohen SE, et al., 2008) in the study to evaluate knowledge of health care provider regarding cardiopulmonary resuscitation of pregnant women. Who pointed out that health care providers knowledge regarding important of basic concepts, including the need for LUD and the potential benefit of early caesarean delivery during cardiac arrest, was inadequate. And added that advanced cardiac life support (ACLS) training is highly recommended.

Concerning effect of educational intervention on maternity nurses knowledge, the present study findings illustrated that was significantly improvement of nurses' knowledge immediately post intervention. These findings were in agreement with (Abd-Allah et al., 2017) in Egypt in the study to investigate effect of educational program on cardiopulmonary resuscitation on nursing student's performance. Who mentioned that educational program is positively improve knowledge of the studied nursing student. In addition the present study findings represented that there was significantly increase of total knowledge score post intervention.

As regard the practice of CPR during pregnancy among maternity nurses, the present study findings proved that the majority of the studied maternity nurses had un satisfactory practice regarding implementation of CPR during pregnancy at pre the program phase, which has been strongly increased immediately post intervention. This result was in agreement with (Elazazay et al., 2012). In Egypt in the study to determine the effect of cardiopulmonary resuscitation training program on nurses knowledge and practice at cancer institute, who mentioned that the nurses performance was very poor in all performance items at pre intervention phase, and their performance has been strongly improved immediately post intervention. These findings may be due to that the use of simulation gives the learner the opportunity to experience uncommon scenarios, receive feedback, and perhaps most importantly correct mistakes before they become part of the learner's. In addition simulation gives

a chance for learners to apply procedure with any fear or distress.

In relation to the studied maternity nurse practice regarding application of chest compression technique, the study finding represented that, nurses had unsatisfactory practice at pre intervention phase and highly improved immediately post intervention. These finding was agreed with (Meissner, 2012). In the study to investigate basic life support skills among high school students before and after resuscitation training who observed significant improvement and good retention rate of CPR practice after program.

Moreover the present study findings showed that the total practice score of the studied nurses regarding CPR during pregnancy was unsatisfactory at pre intervention and become more satisfactory at post intervention, these finding was highly supported by (Elazazay, 2012) in the previous mentioned study, it was indicated that all nurses had unsatisfactory performance at pre the program (100%), while 98.2% of them had good performance immediately post the program.

As regard the relation between the total knowledge score and personal characteristic, the present study findings represented that there were a statistically significant relation between nurses' knowledge and educational qualifications post intervention, this result agrees with (Abd-Allah et al., 2017). In The previous mentioned study, who found a statistically significant relation between level of education and knowledge. Moreover regarding the relation between knowledge and age, the present study proved that there was a no significant relation between knowledge pre, post the program and nurses' age. These finding are in accordance with (Parajulee 2011). in Nepal, in the study to investigate nurses knowledge regarding cardiopulmonary resuscitation in a tertiary care teaching hospital in Nepal, the study findings reported that there was no significant relation between the total knowledge score and age of the respondent, while (Aly, 2010) in Egypt in the study to evaluate impact of a basic life support training program at emergency room on nurses' knowledge and performance, contradicting this result and stated that there is significant relation between age and total basic life support knowledge scores.

As regarding relation between knowledge and years of experience, the finding showed there was no significant relation pre, post the program, this might be attributed that nursing education system is not preparing nurses to be effective in CPR especially and knowledge will be forgotten and deteriorated by time cone in the same line with (Parajulee et al., 2011). In the previous mentioned study, who added that there was no significant relation between the total knowledge score and the duration of experience. On the other hand, the finding was in disagreement with (Al Kandary et al., 2012) in the study to evaluate competence in cardio-pulmonary resuscitation among qualified nurses who proved significant relation between years of experience and knowledge scores.

On other hand there was a highly statically significant relation between total knowledge score of the studied nurses and previous training intervention at the post-intervention phase. This may be due to that training had a pivotal role in improving nurses' knowledge and practice and help nurses to be confident and decrease anxiety toward CPR. These findings agreed with (Kalhori, 2017). In the study to compare the cardiopulmonary resuscitation knowledge between hospital nurses and emergency medical personnel, it was pointed out that there was significantly higher in nurses who had passed basic CPR training courses compared with those who had not ( $P = 0.018$ ).

Furthermore, the results of the current study showed that that there were no statistically significant relations between nurses' practice and educational qualifications pre intervention while there was statically difference at post intervention also regarding the relation between the total practice score and previous training there was a highly statically significant post intervention.

The findings of the study may be related to the previous training which improves nurse's skills as regarding the correlation between total knowledge and practice scores of studied maternity nurse, the present study indicated that there was a highly positive association between total knowledge and practice scores at the post intervention phase, these findings agreed with (Aly, 2010) in the previous mentioned study who reported that there was significant correlation between total basic life support knowledge and practice scores.

## **5. Conclusion**

Study concluded that simulation based intervention is highly improved maternity nurses knowledge, and competence regarding cardio pulmonary resuscitation during pregnancy.

## **6. Recommendation**

Simulation based training regarding cardiopulmonary resuscitation should be provided for all obstetrics health care givers.

## References

- Abd-Allah, E.A., Abd-Alla, K.F., & Mahmoud, A.S. (2017). Effect of Educational Program for Cardiopulmonary Resuscitation Using Simman versus Traditional Manikin on 2nd Year Nursing Students' Performance IOSR. *Journal of Nursing and Health Science*, 6(1), 01-09.
- Adams, J., Jose, R., Lauren, B., Patrick, G., Hughes, M., David, G., ... Michele, L. (2016). Management of Maternal Cardiac Arrest in the Third Trimester of Pregnancy: A Simulation-Based Pilot. *Critical Care Research and Practice*, 2016, 8. <https://doi.org/10.1155/2016/5283765>
- Al Kandary, S., Al Jeheildi, A., Ghayath, T., & Al Haid, N. (2012). Perceived competence in cardio-pulmonary resuscitation, knowledge and practice among qualified nurses in following cardiopulmonary resuscitation training: a review of the literature. *Journal of Advanced Nursing*, 51, 288-297.
- Al-Foudri, H., Kevelighan, E., Catling, S., & CEMACH. (2010). Saving mothers' lives: lessons for anaesthetists, Continuing Education in Anaesthesia. *Crit Care Med.*, 10, 81-87.
- Aly, A. (2010). *Impact of a basic life support training program on nurses' knowledge and performance at emergency room*. Doctorate thesis, Faculty of nursing, Suez Canal University.
- Andreia de, A.R., Nuno C., & Luís, M.G. (2014). Cardiac arrest in pregnancy: best practices are needed. *Acta Obstet Ginecol Port*, 8(2), 164-168.
- Centre for Maternal and Child Enquiries (CMACE). (2006). Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer. The Eighth Report on Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG* 2011, 118(Suppl. 1), 201-203.
- Cohen, S.E., Andes, L.C., & Carvalho, B. (2008). Assessment of knowledge regarding cardiopulmonary resuscitation of pregnant women. *Int J Obstet Anesth*, 17(1), 20-25. <https://doi.org/10.1016/j.ijoa.2007.10.002>
- Damjan, L., Bojan, L., & Jerneja, G. (2012). Impact of additional module training on the level of basic life support knowledge of first year student at the university of Maribor international. *Journal of Emergency Medicine*, 4, 16.
- Demestiha, D., Pantazopoulos, N., & Xanthos, T. (2012). Use of the impedance threshold device in cardiopulmonary resuscitation. *World Journal of Cardiology*, 2(2), 19-26. <https://doi.org/10.4330/wjc.v2.i2.19>
- Einav, S., Matot, I., Berkenstadt, H., et al. (2008). A survey of labour ward clinicians' knowledge of maternal cardiac arrest and resuscitation. *Int J Obstet Anesth*, 17(3), 238-242. <https://doi.org/10.1016/j.ijoa.2008.01.015>
- Elazazay H.M., Abdelazez A.L., & Elsaie, O.A. (2012). Effect of Cardiopulmonary Resuscitation Training Program on Nurses Knowledge and Practice Kuwait. *Bull Alex Fac Med*, 43(2). Hamilton R. Nurses knowledge and skill retention.
- Hui, D., Morrison, L.J., Windrim, R., Lausman, A.Y., et al. (2011). The American Heart Association 2010 guidelines for the management of cardiac arrest in pregnancy: consensus recommendations on implementation strategies. *J. Obstet Gynaecol Can.*, 33, 858-863. [https://doi.org/10.1016/S1701-2163\(16\)34991-X](https://doi.org/10.1016/S1701-2163(16)34991-X)
- Hussain, M., & Lyneham, J. (2009). Cardio-pulmonary resuscitation knowledge among nurses who work in Bahrain. *International Journal of Nursing Practice*, 15(4), 294-302. <https://doi.org/10.1111/j.1440-172X.2009.01752.x>
- Institute of Medicine. (2015). *Strategies to improve cardiac arrest survival: A time to act*. Washington, DC, The National Academies Press.
- Jeejeebhoy, F.M., Zelop, C.M., Lipman, S., et al. (2015). Cardiac Arrest in Pregnancy. *A Scientific Statement From the American Heart Association*. <https://doi.org/10.1161/CIR.0300>
- Lavonas, E.J., Drennan, I.R., Gabrielli, A., et al. (2015). American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care-Part 10: Special Circumstances of Resuscitation. *Circulation*, 132, 501-518. <https://doi.org/10.1161/CIR.0000000000000264>
- Meissner, T., Kloppe, C., & Hanefeld, C. (2012). Basic life support skills of high school students before and after resuscitation training: a longitudinal investigation. *Scandia Journal of Trauma Rhesus*

- EmergencyMedical*, 20, 31. <https://doi.org/10.1186/1757-7241-20-31>
- Micheal, D.D., Pantazopoulos, N., & Xanthos, T. (2014). Use of the impedance threshold device in cardio pulmonary resuscitation. *World Journal of Cardiology*, 2(2), 19-26. <https://doi.org/10.4330/wjc.v2.i2.19>
- Mustafa, M.A., & Jaddoue, B.A. (2014). Assessment of Nurses' Knowledge towards Cardiopulmonary Resuscitation at Al-Najaf City's Teaching Hospital. *Journal of Kufa for Nursing Science*, 4(1).
- Myerburg R.J., & Castellanos. (2015). Cardiac Arrest and Sudden Cardiac Death-Chapter 41(10th ed.). In Bonow R.O., Mann, D.L., Zipes, D.P., & Libby, P. (Eds.), *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. Oxford, UK. Elsevier.
- Nagashima, K., Takahata, O., Fujimoto, A., & Iwasaki, H. (2012). Investigation of nurses' knowledge and experience in cardiopulmonary resuscitation. *Resuscitation*, 14(2), 104-107.
- Neal, J.M., Mulroy, M.F., & Weinberg, G.L. (2012). American Society of Regional Anesthesia and Pain Medicine checklist for managing version. *Reg Anesth Pain Med*, 37, 16-18.
- Parajulee, S., & Selvaraj, V. (2011). Knowledge of nurses towards cardiopulmonary resuscitation in a tertiary care teaching hospital in Nepal. *Journal of Clinical and Diagnostic Research*, 5(8), 1585-1588.
- Reza, P.K., Amir, J., Arsalan, N., Afshin, A., Mohammad, K., Masoud, R., & Mohammad, A. (2017, May-Jun). Assessment of Iranian Nurses and Emergency Medical Personnel in Terms of Cardiopulmonary Resuscitation Knowledge Based on the 2010 Guideline. *Iran J Nurs Midwifery Res.*, 22(3), 184-189.
- Rittenberger, J.C., Kelly, E., Jang, D., & Greer, K.H. (2010). A. Successful outcome utilizing hypothermia after cardiac arrest in pregnancy: a case report. *Crit Care Med*, 36(4), 1354-1356. <https://doi.org/10.1097/CCM.0b013e318169ee99>
- Robertson, L., Schumacher, G., Gosman, R., Kanfer, M.K., & DeVita, M. (2009). Simulation-based crisis team training for multidisciplinary obstetric providers. *Simulation in Healthcare*, 4(2), 77-83. <https://doi.org/10.1097/SIH.0b013e31819171cd>
- Tan, E.K., & Tan, E.L. (2013). Alterations in physiology and anatomy during pregnancy. *Best Pract Res Clin Obstet Gynaecol*, 27, 791-802. <https://doi.org/10.1016/j.bpobgyn.2013.08.001>
- Wong M.K., Morrison L.J., Qiu, F., et al. (2014). Trends in short- and long-term survival among out-of-hospital cardiac arrest patients alive at hospital arrival. *Circulation*, 130(21), 1883-1890. <https://doi.org/10.1161/CIRCULATIONAHA.114.010633>

### Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).