

Effect of Internalized Stigma on Self-esteem and Loneliness among Mentally Ill Patients

Lobna Atya Elwan Ali:-Psychiatric and Mental Health Nursing

Assist. Prof. Dr. Amal Ibrahim Sabra: - Psychiatric and Mental Health Nursing
Department, faculty of Nursing, Tanta University

Dr. Faten Mohammed Ahmed: Psychiatric and Mental Health Nursing
Department, faculty of Nursing, Benha University

Abstract

Recent research studies have shown that, stigma against people with mental illness is universal. Its impact is fundamentally social and can be devastating affects the treatment and prevention of mental disorders. Stigmatized individuals have been found to face a variety of social and emotional consequences including social withdrawal, loss of productivity, lowered self-esteem and increased levels of negative affect. Stigmatization is now recognized as perhaps the central issue facing all who are attempting to understand, prevent and treat mental illness. This study was a descriptive study aiming to evaluate the effect of internalized stigma on self-esteem and loneliness among mentally ill patients. The study was conducted in inpatient ward of Benha mental health hospital. Four instruments were used for collecting the data, socio-demographic and clinical data sheet, Internalized Stigma of Mental Illness Scale, Rosenberg self-esteem scale and UCLA loneliness scale. A convenience sample of 100 psychiatric patients was recruited in this study. The actual study was carried out through three months. The main results revealed that, there were negatively statistical significance relation between internalizing stigma and self-esteem and show positively with loneliness among studied subjects. The study recommended that there is an urgent need for developing strategies to fight internalized stigma associated with mental illness.

Keywords: mental illness - Internalized stigma-self-esteem- loneliness

Introduction

Individuals who are mentally ill are often the targets of bias and stigma and thus susceptible to negative attitudes. As a result, the social and psychological consequences are usually serious and pervasive (*Cheon & Chiao, 2012*). In some studies, stigma toward mental illness has been linked to

poor psychosocial outcomes such as low self-esteem, high rates of depressive symptoms, poor subjective quality of life and loneliness (*Ow & Lee, 2015*) & (*Yen et al., 2015*)

Internalized stigma can create enormous pain for persons with mental illness and may undermine vocational functioning. Additionally it can pervade their lives in many different ways; it diminishes self-esteem and robs people of social opportunities. This can include being denied opportunities such as employment or accommodation because of their illness (*Yanos et al., 2014*).

In fact, stigma is one of the remaining greatest obstacles to the treatment of mental illness. It has been associated with lower self-esteem, depression, feeling misunderstood and ashamed, poor medication adherence, fewer successful social interactions, reduced help seeking, worse recovery, lower quality of life, fewer job opportunities, inadequate health coverage, and fewer leased apartments (*Guyll et al., 2013*).

One of the most tragic consequences of internalized stigma of mental illness is the possibility that it engenders a significant loss of self-esteem specifically, that the stigma of mental illness leads a substantial proportion of people who develop such illnesses to conclude that they are failures or that they have little to be proud of (*Farrelly et al., 2015*).

When anyone, mentally ill or not, does not have enough social contact, it

affects them mentally and even physically. Loneliness creates stress, taking a toll on health (*Hauge & Kirkevold, 2012*). Other things affected can be the ability to learn and memory function. High blood pressure is also seen. It can be the trigger of depression and alcoholism. Imagine the consequences, then, if you are already depressed or have other mental illnesses? Loneliness can make you worse. Loneliness and loss of self-worth lead many mentally ill to believe that they are useless, and so they live with a sense of hopelessness and low self-esteem (*Oexle et al., 2018*).

For this reason, this study was conducted to evaluate the effect of internalized stigma on self-esteem and loneliness among mentally ill patients.

Purpose of the Study

This study aimed to evaluate the effect of internalized stigma on self-esteem and loneliness among mentally ill patients.

Research question:

1-What is the effect of internalized stigma on self-esteem among mentally ill patients?

2- What is the effect of internalized stigma on loneliness among mentally ill patients?

Methods:

Research design:-

Descriptive correlational design was utilized in this study to achieve the aim of the study.

Study setting:-

The study was conducted in psychiatric inpatient ward at psychiatric mental health hospital in Benha city. The hospital is affiliated to General Secretariat of Mental Health. The hospital is divided in three building with capacity of 295 beds and total number of patients 175 patients.

Study Subjects:-

A convenience sample of 100 of psychiatric inpatients was recruited at inpatient psychiatric ward who met the following:-

Inclusion criteria:

- a- The patient age 18:65 years.
- b- Both sexes.
- c-The patient had history of mental illness more than one year.
- d- Hospitalized twice or more.
- e- Willing to participate in the study.

Exclusion criteria:

Patients had the following criteria which itself may be source of stigma:-

- a- Mentally retarded patients.
- b- Physical disability (e.g. hearing or visual disability).
- c- Addiction.
- d- Neurological disorders (e.g. epilepsy).

Instruments:-

Four instruments were used for data collection of the study.

Instrument one: Socio-demographic and clinical data sheet:-

It was developed by the researcher. It was concerned with the following:

- a- Socio-demographic data: It contained 7 items (age, sex, educational level, residence, occupation, marital status and income).
- b- Clinical data: it consisted of 4 items (diagnosis, duration of illness, number of admissions to hospital, mode of current admission).

Instrument two: Internalized stigma of mental illness inventory (ISMI):

It was developed by Ritsher & Jennifer, (2003). It aimed to measure self-stigma of mental illness among persons with mental illness. It was consisting of 29 items,

Effect of Internalized Stigma on Self-esteem and Loneliness among Mentally Ill Patients

each statement was rated on the following 4 point anchored likert scale: 1=strongly disagree, 2=disagree, 3=agree, 4=strongly agree. This scale was divided into 5 subscales: alienation, stereotype endorsement, perceived discrimination, social withdrawal, and stigma resistance.

Subscale 1: The alienation subscale. It consisted of 6 items (1-6) included items such as "having a mental illness has spoiled my life" and "I feel out of place in the world because I have a mental illness".

Subscale 2: The stereotype endorsement subscale. It contained 7 items (7-13) measuring the degree to which respondents agree with common stereotypes about people with mental illness, such as "mentally ill people tend to be violent" and "I can't contribute anything to society because I have a mental illness".

Subscale 3: The discrimination experience subscale. It was composed of 5 items(14-18) intended to capture respondents' perception of the way that they currently tend to be treated by others, such as "people ignore me or take me less seriously just because I have a mental illness" and "people discriminates against me because I have a mental illness".

Subscale 4: The social withdrawal subscale. It was especially heavily

influenced by focus group participants, and contained 6 items(19-24), such as "I don't talk about myself much because I don't want to burden others with my mental illness" and "I avoid getting close to people who don't have mental illness to avoid rejection".

Subscale 5: The stigma resistance subscale. It consisted of 5 items (24-29). It was intended to portray the experience of resisting or being unaffected by internalized stigma, such as "I can have a good, fulfilling life, despite my mental illness", and "I feel comfortable being seen in public with an obviously mentally ill person. The higher score, the higher the internalized stigma patients have.

Instrument three: The Rosenberg Self-esteem scale (Rosenberg, 1965):-

The Rosenberg Self-esteem scale is most widely used self-esteem measure. It was developed by Rosenberg (1965). The scale consists of ten items in which five of them are positive(1,2,4,6,7) and others are negative(3,5,8,9,10). Each item was measured in four point likert-type from strongly agree to strongly disagree.

Scoring system: - to score the items, assign a value to each of ten items as follows: - For items 1, 2, 4, 6, 7 strongly agree = 3 and strongly disagree = 0. For items 3, 5, 8, 9, 10 (which are reversed in valence) strongly agree = 0 and strongly disagree = 3. The

scale ranges from 0- 30 with 30 indicating the highest scale possible.

Low self-esteem: 0-10

Moderate self-esteem: 11-20

High self-esteem: 21-30

Instrument four:-UCLA loneliness scale:-

The UCLA loneliness scale was developed by Russel (1980) to assess subjective feeling of loneliness as well as feeling of social isolation. This scale includes 20 likert-type questions on a four-point likert scale, these 20 item scale ask participants how frequently they agree with statements such as "I feel left out" "I am no longer close to any one" and "my social relationships are superficial". Agreement is measured on a four-point of likert scale ranging from 1=never and 4=often. Items 1-5-6-9-10-15-16-19-20 are reverse score.

Scoring System:

15-20 People attaining this score-range are operating comfortably and experience an average level of loneliness.

21-30 People within this range struggle a little with social interactions, experiencing frequent loneliness.

31-40: Scores falling within this range would indicate a person experiencing severe loneliness.

Methods

Pilot Study

Before actual study a pilot study was conducted on 10% of psychiatric in order to test the clarity of the tools to see if it is clear and fulfilling their purpose. As well as to identify the obstacles and problems that may be encountered during data collection. It served also to estimate the approximate time required for interviewing the patients. After obtaining the result of pilot study, no modifications of tools were done then the final format was developed under the guidance of supervisors. Those patients were excluded later from the actual study.

Ethical consideration

All subjects were informed that participation in the study is voluntary. Anonymity and confidentiality of each participant will be respected and protected. Subjects were informed that the content of the tool will be used for research purpose only. Subjects had the right to refuse to participate in the study or withdrawal at any time. After full explanation of the aim of the study oral consents were taken before participation on the study

Procedure:

- The researcher started data collection by introducing herself to the participant.

Effect of Internalized Stigma on Self-esteem and Loneliness among Mentally Ill Patients

- Oral informed consent was obtained from each participant.
- Brief description of the purpose of the study was given to each participant.
- Data collected were done through interviewing with the patients in hospital in an individual basis. Each interview lasted for 20- 30 minutes, depending on the response of the patients.
- The process of data collection take three months from (July 2017 to September 2017)
- **Statistical analysis**
- The collected data was organized, tabulated, scored, and analyzed using the statistical package for social science (SPSS) version 20. Data was presented using descriptive statistics in the form of frequencies, and percentages and X2 was used mean and stander deviation was used for qualitative data correlation. Confidence spearman was used for correlation. Statistical significant level value was considered when p-value <0.05 and highly significant level value was considered when p-value <0.001 while p-value of >0.05 indicated no significant.

Table (1) shows distribution of the studied patients regarding their socio-demographic characteristics illustrates that about 51% of the studied patients were female. Most of the studied subjects 30% had age from $45 \leq 55$ years. Regarding level of education, 44% of the studied patients read and write and only 1% had university education. On other side 38% were working. The table also revealed that 41% of them were married and 9% were widowed. Finally, more than half of the studied patients (58%) were from rural and their income was enough for 31%, didn't enough for 62% and only 7% can save from it.

Table (2) represents distribution of studied patients regarding their clinical characteristics it was illustrated that duration of illness of only 9% of the studied sample was one year while 36% of them were 3 years and 12% of the studied subjects admitted to psychiatric hospitals for 2-4 times while 45% of them were admitted more than 6 times. It can be noticed that (56%) more than half of the studied subjects were involuntary admission. On the other side more than half of the studied subject's diagnosis was schizophrenia.

Figure (1) describes the distribution of studied patients regarding levels of internalized stigma (ISMI). The results revealed that most of the studied patients (38%, 36%) were suffering from severe and

Results:-

moderate levels of internalized stigma compared to only 13% of subjects suffered from mild and minimal stigma.

Figure (2) reveals distribution of studied patients regarding levels of the Rosenberg self-esteem scale. It showed that the majority of the studied patients had low self-esteem (82%) while only (8%) had high self-esteem.

Figure (3) shows the distribution of the studied subjects regarding levels of UCLA loneliness scale were (44%, 36%) of the studied patients suffered from severe and moderate loneliness and only 20% didn't feel alone.

Table (3) illustrates correlation between internalized stigma (ISMI) and self-esteem among studied patients. It showed that there was highly negative statically correlation between internalized stigma and self-esteem. That means that patients who

had more internalized stigma, become more likely to have low self-esteem.

Table (4) Correlation between internalized stigma of mental illness inventory (ISMI) and UCLA loneliness scale among studied patients is showed in this table. It was clear that there were highly positive significant correlation between internalized stigma and loneliness p-value was 0.000. That means that patients who had more internalized stigma become more likely to have high loneliness score.

Table (1): Distribution of Studied Subjects Regarding Their Socio-Demographic Characteristics

Socio-demographic characteristics items	Studied subjects N=100	
	No	%
Age		
18≤25 years	8	8.0
25≤ 35 years	16	16.0
35≤45 years	25	25.0
45≤ 55 years	30	30.0

Effect of Internalized Stigma on Self-esteem and Loneliness among Mentally Ill Patients

55≤65 years	21	21.0
Sex		
Male	49	49.0
Female	51	51.0
Marital status		
Single	41	41.0
Married	30	30.0
Separated	8	8.0
Divorced	12	12.0
Widowed	9	9.0
Level of education		
Illiterate	31	31.0
Read and write	44	44.0
Intermediate education	24	24.0
University education	1	1.0
Occupation		
Working	38	38.0
Not working	62	62.0
Residence		
Rural	58	58.0
Urban	42	42.0
Income		
Enough	31	31.0
Not enough	62	62.0
Can save from it	7	7.0

Table (2): Distribution of Studied Subjects Sample Regarding Their Clinical Characteristics

Clinical characteristics	Studied subjects N=100	
	No	%
Duration of illness		
1 year	9	9.0

Effect of Internalized Stigma on Self-esteem and Loneliness among Mentally Ill Patients

2 years	25	25.0
3 years	36	36.0
More than 4 years	30	30.0
Number of admission to psychiatric hospitals		
2-3 times	12	12.0
4-6 times	43	43.0
More than 6 times	45	45.0
Mode of current admission		
Voluntary	44	44.0
Involuntary	56	56.0
Diagnosis		
Schizophrenia	59	59.0
Depression	10	10.0
Bipolar disorder	31	31.0




Figure (1): Distribution of studied patients regarding levels of Internalized Stigma (ISMI)




Figure (2): Distribution of Studied Patients Regarding Levels of the Rosenberg Self-Esteem Scale.




Figure (3): Distribution of Studied Subjects Regarding Levels of UCLA Loneliness Scale.

Table (4): Correlation between Internalized Stigma (ISMI) and Self-Esteem among Studied Patients.

Item	Internalized stigma of mental illness inventory (ISMI)

	r	p-value
Rosenberg self-esteem scale	-0.72	0.000

Table (5): Correlation between Internalized Stigma of Mental Illness Inventory (ISMI) and UCLA Loneliness Scale among Studied Patients.

Item	Internalized stigma of mental illness inventory	
	R	p-value
UCLA loneliness scale	0.81	0.000

DISCUSSION:

Mental illnesses are medical conditions that disrupt a person's thinking, feeling, mood, ability to relate to others and daily functioning. Stigmatized individuals have been found to face a variety of social and emotional consequences, including social withdrawal, loss of productivity, lowered self-esteem, and increased levels of negative affect. Stigmatization is now recognized as perhaps the central issue facing all who are attempting to understand, prevent, and treat mental illness (*Hinshaw & Stier, 2008*) & (*Rusch et al., 2014*).

Regarding effect of internalized stigma on self-esteem, it was found that the stigma associated with mental illness harms the self-esteem of many people who have serious mental illnesses. An important consequence of reducing stigma would be to improve the self-esteem of people who have mental illnesses. In the present study, score on the stigma scale were highly negatively correlated with global self-esteem among the studied patients and this result answer our research question that self-stigma has a role on self-esteem among studied participants.

In the Egyptian study carried by *Shalaby et al. (2014)* to investigate" the

role of internalized self-stigma on self-esteem and attitude toward seeking professional psychological help among psychiatric patients", their results indicated that 70% of studied patients have problematic internalized stigma. Also, their study stated that there is a negative relation between internalized self-stigma and self-esteem

Indeed, **Corrigan *et al.* (2012)** stated that the self-esteem of some people with serious psychiatric disorders may be hurt by internalizing stereotypes about mental illness and they also added that people with mental illness face a double-edged sword. Not only do they have to contend with serious, disruptive symptoms, they still have to deal with rampant stigma. Sadly, mental illness is still largely shrouded in stereotypes and misunderstanding.

In this point of the study, **(Bolhari *et al.*, 2002; Geoff, 2011)** mentioned that, a progressive model of self-stigma yields four stages leading to diminished self-esteem and hope: being aware of associated stereotypes, agreeing with them, applying the stereotypes to one's self, and suffering lower self-esteem. They also reported that "People with a mental illness with elevated self-stigma report low self-esteem and low self-image,

and as a result they refrain from taking an active role in various areas of life, such as employment, housing and social life.

Regarding the effect of internalized stigma on loneliness, the findings showed that, most of the studied patients feel loneliness. There was also highly positive statistical significant relation between experiences of internalized stigma and loneliness, this means that a more intense internalized stigma is associated with a higher sense of loneliness; this may be due to feelings of rejection and isolation. The internalized stigma may also lead to depression not only by changing the way people perceive and feel about themselves but also by affecting their perception of their social relations.

Conclusion:

The results of the present study indicated that, internalized stigma played a detrimental role in undermining self-esteem and increasing loneliness among studied subjects.

Recommendations:

Based on the current study findings, the following recommendations are suggested:

- 1- Develop anti-stigma campaigns dealing with mental illness.

- 2- Establishment and empowerment of user organizations is important as well as increasing the awareness of the problem of stigma in professional groups working with the mentally ill.
- 3- Psychological rehabilitative activities should be implemented in psychiatric institutions and rehabilitation centers.
- 4- Planning and implementation of public health awareness programs to raise the orientation toward the nature of psychiatric disorders to minimize the experience of stigma.

References

- Bolhari, J., Nouri ,R., and Ramezani, A. (2002):* In Z. Hussain Khan (Ed.), *Quranic Verses on Mental Health for mental health workers and school counselors* ;p. 22. Tehran: Office for Islamic studies of mental health.
- Corrigan, P., Morris, S., Michaels, P., Rafacz, J. and Rüsçh, N. (2012):* Challenging the public stigma of mental illness a meta-analysis of outcome studies. *Psychiatry Serv*; 63(10):963–973.
- Cheon, B., and Chiao, J. (2012):* Cultural Variation in Implicit Mental Illness Stigma. *Journal of Cross-Cultural Psychology* 43, 1058-1062.
- Farrelly, S., Jeffery, D., and Rüsçh , N. (2015):* The link between mental health-related discrimination and suicidality: service user perspectives. *Psychol Med* 45:2013–2022
- Geoff, J. (2011):* The Relationships of Public and Self-Stigma to Seeking Mental Health Services. *Journal of Mental Health Counseling*, , Vol. 33, *Guyll, M., Spoth, R. and Redmond, C. (2013):* The effects of incentives and research requirements on participation rates for a community-based preventive intervention research study. *Journal Of Primary Prevention* 24, 25-41.
- Hinshaw, P., and Stier, A. (2008):* Stigma as related to mental disorders. *Annual Review of Clinical Psychology*, 4, 367_393. doi: 10.1146 /annurev.clinpsy.4.022007.141245
- Hauge, S. and Kirkevold, M. (2012):* Variations in older persons' descriptions of the burden of loneliness. *Scandinavian Journal of Caring Sciences* 26 (3), 553-560.
- Oexle, N., Rüsçh, N. and Viering, S. (2018):* Self-stigma and suicidality: a longitudinal study. *Eur Arch Psychiatry Clin Neurosci* 267:359–361
- Ow, C .and Lee, B. (2015):* Relationships between perceived stigma, coping orientations, self-esteem, and quality of life in patients with schizophrenia *Asia Pac J Public*

Health.. doi: 10.11 77/
1010539512469246.

spectrum disorders. *Psychiatry
Research* 178 (1), 211–213

Rishter, E. and Jenifer, B. (2003):
Internalized Stigma of Mental
Illness: Psychometric Properties of
a New Measure. *Psychiatry
Research* 121(1):31-49.

*Yen, F., Chen, C., Lee, Y. and Ko, C.
(2015):* Insight and correlates
among outpatients with depressive
disorders. *Psychiatry Services* 56,
599-601

*Rüsch, N., Müller, M., Lay, B. and
Corrigan, W. (2014):* Emotional
reactions to involuntary
psychiatric hospitalization and
stigma-related stress among people
with mental illness. *Eur Arch
Psychiatry Clin Neurosci* 264:35–
43

*Yuksel, C., Bingol, F. and Oflaz,
F. (2013):* Stigma: The cul-de-sac
of the double bind' the perspective
of Turkey; A phenomenological
study. *Journal of Psychiatric and
Mental Health Nursing*, 21, 667–
678.

Rosenberg, M. (1965): Society and the
adolescent self-Image. Princeton,
NJ:Princeton University Press.
*Journal of Personality and Social
Psychology*.89, 623-642.

Russel, L. (1980): Concurrent and
discriminant validity evidence.
*Journal of Personality and Social
Psychology*. 39.472.480.

*Shalaby, M., Sabra, A., and Mohamed, S.
(2014):* The Role Of Internalized
Self Stigma On Self Esteem And
Attitude Toward Seeking
Professional Psychological Help
Among Psychiatric Patients.
*International Journal of Current
Research, Vol. 6, Issue,
11,pp.10162-10169, November,
2014.*

Yanos, P., Lysaker, P. and Roe, D. (2014):
Internalized stigma as a Barrier to
Improvement Prevalence in
vocational functioning among
people with schizophrenia